



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

FORMATION OF SPORANGIA IN STEMONITIS

Hilton (Jour. Q. M. C., Apr. 1916) contributes an interesting note on the method of forming sporangia in this common mycetozoon. He had the good fortune to find considerable masses of plasmodium just in condition to observe the whole process. At noon the plasmodium formed somewhat rounded, solid, cushion-shaped masses. This surface differentiated in about a quarter of an hour into frothy, bubble-like hemispheres which divided and covered the entire surface regularly. By 4:00 P. M. each mass of hemispheres had contracted in width and increased in height; and the basal part constricted into flutings corresponding to the surface hemispheres. These flutings gradually contracted to pillars, the creamy protoplasm withdrawing more and more to the upper third. In another half hour black stalks were visible as the cores of the pillars. By 8:30 o'clock all the protoplasm had risen clear of the substratum and the still cohering heads of the sporangia appeared resting on a forest of black stalks. By 10:00 P. M. the sporangia had virtually assumed their permanent shape and were beginning to darken.

A DROUTH-ENDURING ZYGNEMA

Fritsch (Ann. Bot. 1916, pp. 135-149) reports upon a *Zygnema* especially adapted to terrestrial conditions. The longitudinal walls become much thickened, showing two or three successive layers. The outer layer of the wall is mucilaginous. This doubtless is an adaptation which prevents too rapid drying and aids resorption of water on return of moist conditions. It was found on Hindhead Common (England.)

There are two chloroplasts in the mature cell. Division takes place by an infolding or growth from the inner layer of the cell wall. The growth of this gradually constricts the protoplast, but the division may not be completed for considerable time. In this way the two daughter protoplasts have at first a connection thru the center of this plate.

With the oncoming of drouth the fat globules of the cell pass to the outer surface, and form there a dense layer just beneath the cell wall. When the plant begins to absorb water the fat drops